arXiv:1504.01378v2 [cond-mat.stat-mech] 29 Dec 2016

The effects of the chemical potential in a BE distribution and the fractional parameter in a distribution with Mittag–Leffler function

Minoru Biyajima[a], Takuya Mizoguchi[b], Naomichi Suzuki[c]

[a]*Department of Physics, Shinshu University, Matsumoto 390-8621, Japan*
[b]*National Institute of Technology, Toba College, Toba 517-8501, Japan*
[c]*Matsumoto University, Matsumoto 390-1295, Japan*

Abstract

The fractional Planck distribution is calculated by applying the Caputo fractional derivative with order p ($p > 0$) to the equation proposed by Planck in 1900. In addition, the integral representation of the Mittag–Leffler function is employed to obtain a new formula for the fractional BE distribution, which is then used to analyze the NASA COBE monopole data. Based on this analysis, an identity $p \simeq e^{-\mu}$ is found, where μ is the dimensionless constant chemical potential that was introduced to the BE distribution by the NASA COBE collaboration.

Keywords: Planck distribution; fractional calculus; Mittag–Leffler function; Integral representation of ML function; Bose–Einstein distribution; NASA COBE data

---

1. Introduction

There has been recent progress in physical sciences studies [1, 2, 3, 4, 5, 6] that are based on fractional calculus [7, 8, 9, 10, 11, 12, 13]. As part of these contributions, we investigate the NASA COBE monopole data [14, 15, 16] by utilizing the Bose–Einstein (BE) distribution and a fractional calculus based distribution [17, 18, 6]. A well-known solution for Kompaneets equation [19, 20], which describes the photons distribution in the early Universe, is given by:

$$
\begin{aligned}
U^{(\mathrm{BE})}(T,\ \nu,\ \mu) &= \frac{C_B}{e^{x+\mu}-1} \xrightarrow{|\mu|\ll 1} C_B\left[\frac{1}{e^x-1} - \mu\frac{e^x}{(e^x-1)^2}\right] \\
&= F^{(\mathrm{PD})} + F_2^{(\mathrm{BE})}(x,\ \mu), \qquad (1)
\end{aligned}
$$

where $x = h\nu/k_BT$ and $C_B = 2h\nu^3/c^2$. μ is a dimensionless chemical potential. See Table 1 I) Kompaneets equation.

On the other hand, the fractional calculus based photons distribution of the Universe is given by (See Table 1 II) Fractional calculus and III) Planck distribution in 1900 [21, 22, 23]):

$$U(x) = \frac{C_B}{E_p(x^p) - 1}, \qquad (2)$$

where p is the fractional parameter. $E_p(x^p)$ is the Mittag–Leffler (ML) function defined as

$$E_p(x^p) = \sum_{n=0}^{\infty} \frac{x^{np}}{\Gamma(np+1)}. \qquad (3)$$

Email addresses: `biyajima@azusa.shinshu-u.ac.jp` (Minoru Biyajima), `mizoguti@toba-cmt.ac.jp` (Takuya Mizoguchi), `suzuki@matsu.ac.jp` (Naomichi Suzuki)

The following digamma function ψ is used in ref. [6] to analyze the NASA COBE data,

$$\begin{aligned} \text{Eq. (2)} \xrightarrow{|p-1|\ll 1} \quad & C_B\left[\frac{1}{e^x-1}+\frac{p-1}{(e^x-1)^2}\sum_{k=0}^{\infty}\frac{kx^k[\psi(k+1)-\ln x]}{\Gamma(k+1)}\right] \\ & = F^{(\mathrm{PD})}+F_2^{(\mathrm{FC})}(x,\ p-1), \end{aligned} \tag{4}$$

where $\psi(z) = d(\ln\Gamma(z))/dz$.

The COBE data is analyzed using Eqs. (1) and (2), thereby yielding the following estimated values:

$$\begin{aligned} & T = 1/k_B\beta = 2.725\ \mathrm{K},\ |\mu| < 7.58\times 10^{-5}, \\ & |p-1| < 8.09{\times}10^{-5},\ |\mu|{\times}I_1 < 5.47{\times}10^{-4}\ (1.00), \\ & |p-1|\times I_2 < 5.60\times 10^{-4}\ (1.03), \end{aligned} \tag{5}$$

where, with Riemann's ζ function $\zeta(3)$,

$$\begin{aligned} I_1 &= \int_0^{\infty}\frac{x^3\cdot e^x}{(e^x-1)^2}dx = 3\cdot 2\cdot\zeta(3), \\ I_2 &= \int_0^{\infty}\frac{x^3}{(e^x-1)^2}\sum_{k=0}^{\infty}\frac{kx^k[\psi(k+1)-\ln x]}{\Gamma(k+1)}dx \\ &= \sum_{k=0}^{\infty}\sum_{m=2}^{\infty}\frac{k(m-1)}{\Gamma(k+1)}\left(\frac{1}{m}\right)^{k+4}\Gamma(k+4)\left[\ln m-\left(\frac{1}{k+3}+\frac{1}{k+2}+\frac{1}{k+1}\right)\right]. \end{aligned}$$

We would like to pay our attention to these similar values, μ and $p-1$. The ratio between $|\mu|\times I_1$ and $|p-1|\times I_2$ is 1.00 to 1.03.

The same analysis of the COBE data is presented in this study, where we would like to adopt an integral representation of the ML function:

$$E_p(x^p) = \frac{e^x}{p} + \delta(p,\ x), \tag{6}$$

where

$$\delta(p,\ x) = -\frac{\sin(p\pi)}{\pi}\int_0^{+\infty}\frac{y^{p-1}e^{-xy}}{y^{2p}-2y^p\cos(p\pi)+1}dy. \tag{7}$$

A detailed derivation of the integral representation is supplied in §2. The magnitude of the integral representationi$\delta(x,\ p)$) contribution is estimated through concrete analysis of the COBE data in §3. Through such analysis of the COBE data, the following relation ensues:

$$\mu = -\ln p\,. \tag{8}$$

The concluding remarks and discussion are provided in §4.

2. Integral representation of the Mittag–Leffler function

The integral representation of the Mittag–Leffler (ML) function $E_\alpha(z)$ [11, 9, 12, 13] is given by,

$$E_\alpha(z) = -\frac{1}{2\pi i\alpha}\int_{\gamma(\varepsilon;\delta)}\frac{e^{\zeta^{1/\alpha}}}{\zeta-z}d\zeta + \frac{1}{\alpha}e^{z^{1/\alpha}}, \quad z\in G^{(+)}(\varepsilon;\delta), \tag{9}$$

$$E_\alpha(z) = -\frac{1}{2\pi i\alpha}\int_{\gamma(\varepsilon;\delta)}\frac{e^{\zeta^{1/\alpha}}}{\zeta-z}d\zeta, \quad z\in G^{(-)}(\varepsilon;\delta), \tag{10}$$

Table 1: Kompaneets equation, the fractional and the basic equations of the Planck distribution.

I) Kompaneets Eq. describing the photons distributions in the early Universe [19, 20] It is given by $\partial f/\partial t = C_\kappa x_e^{-2} \frac{\partial}{\partial x_e} x_e^4 (\partial f/\partial x_e + f + f^2)$ where $C_\kappa = (k_B T_e/m_e c^2)(n_e \sigma_e/c)$ and $x_e = h\nu/k_B T_e$. T_e, T, n_e, and σ_e are the electron temperature, radiation temperature, electron density, and Thomson's cross-section, respectively. As a stationary solution, we derive the Bose –Einstein (BE) distribution with the chemical potential, $f(x) = 1/(ce^x - 1) = 1/(e^{x_e+\mu} - 1)$.
II) Fractional calculus [7] For the stationary solution in I) and a solution in III), if an inverse function $R(x) = 1/f(x)$ exists, the following equation is obtained, $\partial R/\partial x = R(x) + 1$. The Caputo derivative in fractional calculus is applied to the previous equation, *i.e.*, ${}_0 D_x^p R(x) = R(x) + 1$, The Caputo fractional derivative of the $f(x)$ function for $m = 1, 2, \ldots$ is given by ${}_0^C D_x^p f(x) = \frac{1}{\Gamma(m-p)} \int_0^x (x-\tau)^{m-p-1} f^{(m)}(\tau) d\tau, \quad (m-1 < p < m)$, and $\lim_{p \to m} {}_0^C D_x^p f(x) = f^{(m)}(x) = d^m f(x)/dx^m$. The following distribution is calculated using the ML function, $f(x) = 1/R(x) = 1/(E_p(x^p) - 1)$.
III) Planck distribution in 1900 [21, 22, 23, 24] Planck utilized the following equation to describe the photons distribution U, $\partial U/\partial \beta = -U - U^2$, Adopting the aforementioned method in section II), the same expression is deduced.

in the complex plane, $z = x + iy$ ($x = Re\, z, y = Im\, z$), under the conditions,

$$0 < \alpha < 2, \quad \pi\alpha/2 < \delta \leq \min\{\pi, \pi\alpha\}. \tag{11}$$

As shown in Fig. 1, the contour $\gamma(\varepsilon; \delta)$ ($\varepsilon > 0$, $0 < \delta \leq \pi$) is comprised of the following three parts: (i) ray $S_{-\delta}$ ($\arg \tau = -\delta$, $|\tau| \geq \varepsilon$), (ii) circular arc $C_\delta(0; \varepsilon)$ ($-\delta \leq \arg \tau \leq \delta$, $|\tau| = \varepsilon$), (iii) ray S_δ ($\arg \tau = \delta$, $|\tau| \geq \varepsilon$). The left hand side of the contour $\gamma(\varepsilon; \delta)$, where the origin O is included, and is denoted by $G^{(-)}(\varepsilon; \delta)$, while the other side is denoted by $G^{(+)}(\varepsilon; \delta)$D

After integrating along the contour $\gamma(\varepsilon; \delta)$ while taking the $\varepsilon \to 0$ limit into consideration, Eqs. (9) and (10) respectively reduce to,

$$E_\alpha(z) = -\frac{z \sin(\pi\alpha)}{\pi\alpha} \int_0^{+\infty} \frac{e^{-r^{1/\alpha}}}{r^2 - 2rz\cos(\pi\alpha) + z^2} dr + \frac{1}{\alpha} e^{z^{1/\alpha}}, \quad \in G^{(+)}(+0; \delta), \tag{12}$$

$$E_\alpha(z) = -\frac{z \sin(\pi\alpha)}{\pi\alpha} \int_0^{+\infty} \frac{e^{-r^{1/\alpha}}}{r^2 - 2rz\cos(\pi\alpha) + z^2} dr, \quad z \in G^{(-)}(+0; \delta). \tag{13}$$

If $0 < \alpha \leq 1$, the explicit expression for $E_\alpha(z)$ is obtained through Eqs.(12) and (13). If $\alpha > 1$, using the ML function's summation formula,

$$E_\alpha(z) = \frac{1}{m} \sum_{h=0}^{m-1} E_{\alpha/m}(z^{1/m} e^{i2\pi h/m}), \tag{14}$$

we can express $E_\alpha(z)$ through those with suffixes $\alpha/m \leq 1$. Then, the ML function's expression is given by the procedure in the case of $0 < \alpha < 1$.



Figure 1: The contour $\gamma(\varepsilon;\delta)$ composed of the ray $S_{-\delta}$, circular arc $C_\delta(0;\varepsilon)$ and ray S_δ

In Fig. 2, we numerically examine the behavior of $\delta(p,\ x)$ in the ML function $E_p(x^p)$ for $0 < p < 2$ and $x \geq 0$. The integral representation $\delta(p,\ x)$ for $0 < p$ defined in Eq.(7) approaches to zero in the limit of $x \to \infty$. Function $\delta(p,\ x)$ at fixed p for $0 < p < 1$ is negative and increases monotonously with x, and $\delta(p,\ x)$ for $1 < p < 2$ is positive and decreases monotonously with x. For $0 < p < 2$, the following limit can be derived through analytic calculations [25]:

$$|\delta(p,\ x)| \leq |\delta(p,\ 0)| = |p-1|/p. \tag{15}$$



Figure 2: a) Contour maps of $\delta(p,\ x)$ = constant for $0.5 < p < 1$. b) Contour maps of $\delta(p,\ x)$ = constant for $1 < p < 2$.

3. Analysis of COBE data by Eqs. (2) and (6)

Expanding Eq. (2), providing that $(p-1) \ll 1$, as follows

$$U(T,\nu,p) \quad = \quad \frac{C_B}{e^x-1} + \frac{C_B e^x \ln p}{(e^x-1)^2} - \frac{C_B\delta(p,\ x)}{(e^x-1)^2}$$

$$= \quad F^{(\mathrm{PD})}(x) + F_2^{(\mathrm{FC})}(x,\, \ln p) - F_3^{(\mathrm{FC})}(x,\, \delta(p,\, x)), \tag{16}$$

we can analyze the NASA COBE data in Fig. 3. For comparison, the results on $F_2^{(\mathrm{FC})}(x,\ \ln p)$ and $F_3^{(\mathrm{FC})}(x,\ \delta(p,\ x))$



Figure 3: Analysis of the COBE monopole data by Eqs. (16), (1) and (4). a) Eq. (16). b) Eqs. (1), (4) and (16). The magnitudes of error bars mean that 400 times real error bars.

are shown separately. As is seen in Fig. 3, it can be said that contribution from $F^{(\mathrm{PD})}(x)$ is much bigger than that of $F_3^{(\mathrm{FC})}(x,\ \delta(p,\ x))$. Then we can directly compare Eq. (1) with e^x/p and obtain Eq. (8): $\mu = -\ln p$. From this expression, p can be regarded as an inverse fugacity [26, 27], provided that NASA COBE data are used and the magnitude of $\delta(p,\ x)$ is very small.

Analysis of NASA COBE data by Eq. (16) is presented in Table 2.

Table 2: Numerical proof of equivalences between Eqs. (2) and (16) is shown at typical observed values of NASA COBE data. Our result of analysis by Eq. (16) is as follows: $T = 2.725008 \pm 0.000026$ K, $|p-1| = 8.09 \times 10^{-5}$ and χ^2/NDF = 45.01/41. It should be noticed that we obtain the following values, $T = 2.725016 \pm 0.000008$ K and χ^2/NDF = 45.10/42, provided that the first term $F^{(\mathrm{PD})}$ is only utilized.

ν/c [cm^{-1}]	4.08	8.62	13.16
Eq. (2)	354.0636052	271.4238381	87.02961125
Eq. (16)	354.0636052	271.4238381	87.02961124
$F_1^{(\mathrm{PD})}$	354.0594726	271.4209229	87.02868402
$F_2^{(\mathrm{FC})}$	$0.4.263569862\times10^{-3}$	2.920132007×10^{-3}	9.273229351×10^{-4}
$F_3^{(\mathrm{FC})}$	1.310610486×10^{-4}	4.892780012×10^{-6}	1.016807344×10^{-7}

4. Concluding remarks and discussion

C1) We have shown that the ML function of Eq. (6) is decomposed into two functions e^x/p and $\delta(p,\ x)$: The various properties of the integral representation $\delta(p,\ x)$ are investigated. In particular, through the analysis of the COBE monopole data, it is shown that the magnitude of $\delta(p,\ x)$ is much smaller than that of the Planck distribution. See, Fig. 3a).

C2) In our previous paper [6], we observed $|\mu| = |p-1|$ through the numerical analysis of the COBE monopole data by Eqs. (1), (2) and (4). In the present paper, we have obtained Eq. (8) i.e., $\mu = -\ln p$ in the analytic form.

C3) Combining the analytic relation of Eq. (8) with the numerical results of Table 2, the following relation is obtained,

$$|\mu| = \ln(|p-1| + 1). \tag{17}$$

D1) Using the present study, the interesting result given by Eq. (17) is achieved. In the near future, the effect of the fractional parameter on other fields (BE condensation) should be investigated. From Eqs. (2) and (6), we obtain the following formula,

$$g_\nu(p) = \frac{1}{\Gamma(\nu)} \int_0^\infty \frac{x^{\nu-1}}{p^{-1}e^x - 1} dx = \sum_{l=1}^{\infty} \frac{p^l}{l^\nu},\ (p < 1), \tag{18}$$

It is well-known that this formula is given in Refs. [26, 27, 28]

Acknowledgements

We would like to acknowledge Prof. M. Caputo for his kindness. Moreover, one of authors (M.B.) would like to thank the Department of Physics at Shinshu University for their hospitality.

References

[1] R. Metzler and J. Klafter, Phys. Rep. **339**, 1 (2000).

[2] B. J. West, M. Bologna and P. Grigolini, *Physics of Fractal Operators*, (Springer-Verlag, 2003).

[3] R. Hilfer, J. Phys. Chem. **143**, 3914 (2000).

[4] E. Barkai, Phys. Rev. E **63**, 046118 (2001).

[5] N. Suzuki and M. Biyajima, Phys. Rev. E **65**, 016123 (2001).

[6] M. Biyajima, T. Mizoguchi and N. Suzuki, Physica **A 440**, 129 (2015).

[7] M. Caputo, Geophys. J. R. Astr. Soc. **13**, 529 (1967); M. Caputo and F. Mainardi, Riv. Nuov. Cim. (Ser. II), **1**, 161 (1971); M. Caputo, J. M. Carcione and M. A. B. Botelho, Fract. Calc. Appl. Anal. **18**, 3 (2015).

[8] B. Ross (Ed.), *Fractional Calculus and Its Applications*, Lecture Notes in Mathematics, vol. 457 (Springer-Verlag, New York, 1975).

[9] I. Podlubny, *Fractional Differential Equations* (Academic Press, San Diego, 1999).

[10] F. Mainardi, Y. Luchiko and G. Panini, Frac. Calc. and Appl. Anal. **4**, 153 (2001).

[11] M. M. Dzherbashian, *Harmonic analysis and Boundary Value Problems in the Complex Domain*, (Birkhauser, Basel, 1993).

[12] R. Gorenflo, J. Loutchko and Y. Luchko, Frac. Calc. Appl. Anal. **5**, 491 (2002).

[13] R. Gorenflo, A. A. Kilbas, F. Mainardi and S. V. Rogosin, *Mittag-Leffler Functions, Related Topics and Applications* , (Springer, Heiderberg, 2014).

[14] J.C. Mather et al., Astrophys. **J. 420**, 439 (1994); See also, D.J. Fixsen et al., Astrophys. **J. 473**, 576 (1996); D.J. Fixsen and J.C. Mather, Astrophys. **J. 581**, 817 (2002).

[15] COBE/FIRAS CMB monopole spectrum, May 2005,
http://lambda.gsfc.nasa.gov/product/cobe/firas_monopole
_get.cfm ;

[16] R.Durrer, *The Cosmic Microwave Background*, (Cambridge University press, 2008).

[17] H. Ertik et al., Physica **A 388**, 4573 (2009).

[18] M. Biyajima and T. Mizoguchi, Phys. Lett. **A 376** (2012) 3567; See also, M. Biyajima and T. Mizoguchi, Astrophys. Space Sci. **350**, 317 (2014).

[19] A. S. Kompaneets, Sov. Phy. JETP **4**, 730 (1957).

[20] R. Weymann, Phys. Fluids **8**, 2112 (1965).

[21] M. Planck, *Ueber irreversible Strahlungsvorgänge*, Ann. d. Phys. **1**, 69 (1900).

[22] M. Planck, Verh. Deutsch. Phys. Ges. **2**, 202 and 237 (1900).

[23] A. Sommerfeld, *Thermodynamics and Statistical Mechanics* (Academic Press, Yew York, 1956).

[24] C. Tsallis, *Introduction to Nonextensive Statistical Mechanics* (Springer Science+Business media LLC, 2009).

[25] I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series and Products*, (Academic Press, Yew York, 1965).

[26] R. K. Pathria, *Statistical Mechanics*, (Second Ed., Elsevier, 1996).

[27] K. Huang, *Statistical Mechanics*, (Second ed., John Wiley & Suns, Yew York, 1987).

[28] C. J. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases*, (Cambridge University Press, 2002).